

TRANSLATOR'S DECLARATION

I, Janet Hope, BSc(Hons.), MIL., MITI., translator to Messrs. Taylor and Meyer of 20 Kingsmead Road, London, SW2 3JD, Great Britain, verify that I know well both the German and the English language, that I have prepared the attached English translation of 33 pages of a German Patent application in the German language with the title:

Neue für das cstA-Gen kodierende Nukleotidsequenzen

identified by the code number 000173 BT at the upper left of each page and that the attached English translation of this document is a true and correct translation of the document attached thereto to the best of my knowledge and belief.

I further declare that all statements made of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements are made with the knowledge that wilful false statements and the like are punishable by fine or imprisonment, or both, under 18 USC 1001, and that such false statements may jeopardize the validity of this document.

Signed:



Dated: 13th November 2003

32301.195

FEDERAL REPUBLIC OF GERMANY

Certificate of Priority for Filing of a Patent Application

Filing number: 100 42 051.6

Filing date: 26th August 2000

Applicant/Proprietor: Degussa AG, Düsseldorf/Germany

First applicant: Degussa-Hüls AG,
Frankfurt am Main/Germany

Title: New nucleotide sequences which code for the cstA
gene

IPC: C 07 H, C 12 N, C 12 Q

**The attached papers are a true and accurate reproduction of the original
documents for this patent application.**

Munich, 5th July 2001

**On behalf of the President of the German
Patent and Trade Mark Office**

(signature)

Sieck

32301.195

New nucleotide sequences which code for the cstA gene

The invention provides nucleotide sequences from coryneform bacteria which code for the cstA gene and a process for the fermentative preparation of amino acids, in particular L-lysine, using bacteria in which the cstA gene is enhanced.

Prior art

L-Amino acids, in particular L-lysine, are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal nutrition.

It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular *Corynebacterium glutamicum*. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the process can relate to fermentation measures, such as, for example, stirring and supply of oxygen, or the composition of the nutrient media, such as, for example, the sugar concentration during the fermentation, or the working up to the product form by, for example, ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites or are auxotrophic for metabolites of regulatory importance and produce amino acids are obtained in this manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of *Corynebacterium* strains which produce L-amino acid, by amplifying individual amino acid biosynthesis genes and investigating the effect on the amino acid production.

Object of the invention

Where L-amino acids or amino acids are mentioned in the following, this means one or more amino acids, including their salts, chosen from the group consisting of L-

5 asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine.

The inventors had the object of providing new measures for
10 improved fermentative preparation of amino acids, in particular L-lysine.

Description of the invention

When lysine or L-lysine are mentioned in the following, not only the base but also the salts, such as e.g. lysine
15 monohydrochloride or lysine sulfate, are meant by this.

The invention provides an isolated polynucleotide from coryneform bacteria, comprising a polynucleotide sequence which codes for the cstA gene, chosen from the group consisting of

- 20 a) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2,
- b) polynucleotide which codes for a polypeptide which
25 comprises an amino acid sequence which is identical to the extent of at least 70% to the amino acid sequence of SEQ ID No. 2,
- c) polynucleotide which is complementary to the polynucleotides of a) or b), and

d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequence of a), b) or c),

5 the polypeptide preferably having the activity of carbon starvation protein A.

The invention also provides the abovementioned polynucleotide, this preferably being a DNA which is capable of replication, comprising:

- (i) the nucleotide sequence shown in SEQ ID No. 1, or
- 10 (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
- (iii) at least one sequence which hybridizes with the sequence complementary to sequence (i) or (ii),
- 15 and optionally
- (iv) sense mutations of neutral function in (i).

The invention also provides

a polynucleotide comprising the nucleotide sequence as shown in SEQ ID No. 1;

20 a polynucleotide which codes for a polypeptide which comprises the amino acid sequence as shown in SEQ ID No. 2;

a vector containing the polynucleotide according to the invention, in particular a shuttle vector or plasmid
25 vector, and

coryneform bacteria serving as the host cell, which contain the vector or in which the *cstA* gene is enhanced.

The invention also provides polynucleotides which substantially comprise a polynucleotide sequence, which are

obtainable by screening by means of hybridization of a corresponding gene library of a coryneform bacterium, which comprises the complete gene or parts thereof, with a probe which comprises the sequence of the polynucleotide

- 5 according to the invention according to SEQ ID No.1 or a fragment thereof, and isolation of the polynucleotide sequence mentioned.

Polynucleotide sequences according to the invention are suitable as hybridization probes for RNA, cDNA and DNA, in
10 order to isolate, in the full length, nucleic acids or polynucleotides or genes which code for carbon starvation protein A, or to isolate those nucleic acids or polynucleotides or genes which have a high similarity of sequence with that of the cstA gene.

- 15 Polynucleotide sequences according to the invention are furthermore suitable as primers with the aid of which DNA of genes which code for carbon starvation protein A can be prepared with the polymerase chain reaction (PCR).

Such oligonucleotides which serve as probes or primers
20 comprise at least 30, preferably at least 20, very particularly preferably at least 15 successive nucleotides. Oligonucleotides which have a length of at least 40 or 50 nucleotides are also suitable.

"Isolated" means separated out of its natural environment.

- 25 "Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these to be non-modified RNA or DNA or modified RNA or DNA.

The polynucleotides according to the invention include a polynucleotide according to SEQ ID No. 1 or a fragment
30 prepared therefrom and also those which are at least 70%, preferably at least 80% and in particular at least 90% to 95% identical to the polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom.

"Polypeptides" are understood as meaning peptides or proteins which comprise two or more amino acids bonded via peptide bonds.

The polypeptides according to the invention include a
5 polypeptide according to SEQ ID No. 2, in particular those with the biological activity of carbon starvation protein A, and also those which are at least 70%, preferably at least 80% and in particular which are at least 90% to 95% identical to the polypeptide according to SEQ ID No. 2 and
10 have the activity mentioned.

The invention moreover provides a process for the fermentative preparation of amino acids, in particular L-lysine, using coryneform bacteria which in particular already produce amino acids, and in which the nucleotide
15 sequences which code for the *cstA* gene are enhanced, in particular over-expressed.

The term "enhancement" in this connection describes the increase in the intracellular activity of one or more enzymes in a microorganism which are coded by the
20 corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or using a gene which codes for a corresponding enzyme having a high activity, and optionally combining these measures.

The microorganisms which the present invention provides can
25 prepare L-amino acids, in particular L-lysine, from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They can be representatives of coryneform bacteria, in particular of the genus *Corynebacterium*. Of the genus *Corynebacterium*,
30 there may be mentioned in particular the species *Corynebacterium glutamicum*, which is known among experts for its ability to produce L-amino acids.

Suitable strains of the genus *Corynebacterium*, in particular of the species *Corynebacterium glutamicum* (*C. glutamicum*), are in particular the known wild-type strains

- 5 *Corynebacterium glutamicum* ATCC13032
- Corynebacterium acetoglutamicum* ATCC15806
- Corynebacterium acetoacidophilum* ATCC13870
- Corynebacterium thermoaminogenes* FERM BP-1539
- Corynebacterium melassecola* ATCC17965
- Brevibacterium flavum* ATCC14067
- 10 *Brevibacterium lactofermentum* ATCC13869 and
- Brevibacterium divaricatum* ATCC14020

and L-lysine-producing mutants or strains prepared therefrom, such as, for example

- 15 *Corynebacterium glutamicum* FERM-P 1709
- Brevibacterium flavum* FERM-P 1708
- Brevibacterium lactofermentum* FERM-P 1712
- Corynebacterium glutamicum* FERM-P 6463
- Corynebacterium glutamicum* FERM-P 6464 and
- Corynebacterium glutamicum* DSM5715.

- 20 The inventors have succeeded in isolating the new *cstA* gene of *C. glutamicum* which codes for carbon starvation protein A.

- To isolate the *cstA* gene or also other genes of *C. glutamicum*, a gene library of this microorganism is first
- 25 set up in *Escherichia coli* (*E. coli*). The setting up of gene libraries is described in generally known textbooks and handbooks. The textbook by Winnacker: *Gene und Klone, Eine Einführung in die Gentechnologie* (Verlag Chemie, Weinheim, Germany, 1990), or the handbook by Sambrook et
 - 30 al.: *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) may be mentioned as an example. A well-known gene library is that of the *E. coli* K-12 strain W3110 set up in λ vectors by Kohara et al. (Cell 50, 495 -508 (1987)). Bathe et al. (Molecular and

General Genetics, 252:255-265, 1996) describe a gene library of *C. glutamicum* ATCC13032, which was set up with the aid of the cosmid vector SuperCos I (Wahl et al., 1987, Proceedings of the National Academy of Sciences USA, 84:2160-2164) in the *E. coli* K-12 strain NM554 (Raleigh et al., 1988, Nucleic Acids Research 16:1563-1575).

Börmann et al. (Molecular Microbiology 6(3), 317-326 (1992)) in turn describe a gene library of *C. glutamicum* ATCC13032 using the cosmid pH79 (Hohn and Collins, Gene 11, 291-298 (1980)). To prepare a gene library of *C. glutamicum* in *E. coli* it is also possible to use plasmids such as pBR322 (Bolivar, Life Sciences, 25, 807-818 (1979)) or pUC9 (Vieira et al., 1982, Gene, 19:259-268). Suitable hosts are, in particular, those *E. coli* strains which are restriction- and recombination-defective. An example of these is the strain DH5 α mc^r, which has been described by Grant et al. (Proceedings of the National Academy of Sciences USA, 87 (1990) 4645-4649). The long DNA fragments cloned with the aid of cosmids can in turn be subcloned in the usual vectors suitable for sequencing and then sequenced, as is described e.g. by Sanger et al. (Proceedings of the National Academy of Sciences of the United States of America, 74:5463-5467, 1977).

The resulting DNA sequences can then be investigated with known algorithms or sequence analysis programs, such as e.g. that of Staden (Nucleic Acids Research 14, 217-232(1986)), that of Marck (Nucleic Acids Research 16, 1829-1836 (1988)) or the GCG program of Butler (Methods of Biochemical Analysis 39, 74-97 (1998)).

The new DNA sequence of *C. glutamicum* which codes for the *cstA* gene and which, as SEQ ID No. 1, is a constituent of the present invention has been obtained in this manner. The amino acid sequence of the corresponding protein has furthermore been derived from the present DNA sequence by

the methods described above. The resulting amino acid sequence of the cstA gene product is shown in SEQ ID No. 2.

Coding DNA sequences which result from SEQ ID No. 1 by the degeneracy of the genetic code are also a constituent of the invention. In the same way, DNA sequences which hybridize with SEQ ID No. 1 or parts of SEQ ID No. 1 are a constituent of the invention. Conservative amino acid exchanges, such as e.g. exchange of glycine for alanine or of aspartic acid for glutamic acid in proteins, are furthermore known among experts as "sense mutations" which do not lead to a fundamental change in the activity of the protein, i.e. are of neutral function. It is furthermore known that changes on the N and/or C terminus of a protein cannot substantially impair or can even stabilize the function thereof. Information in this context can be found by the expert, inter alia, in Ben-Bassat et al. (Journal of Bacteriology 169:751-757 (1987)), in O'Regan et al. (Gene 77:237-251 (1989)), in Sahin-Toth et al. (Protein Sciences 3:240-247 (1994)), in Hochuli et al. (Bio/Technology 6:1321-1325 (1988)) and in known textbooks of genetics and molecular biology. Amino acid sequences which result in a corresponding manner from SEQ ID No. 2 are also a constituent of the invention.

In the same way, DNA sequences which hybridize with SEQ ID No. 1 or parts of SEQ ID No. 1 are a constituent of the invention. Finally, DNA sequences which are prepared by the polymerase chain reaction (PCR) using primers which result from SEQ ID No. 1 are a constituent of the invention. Such oligonucleotides typically have a length of at least 15 nucleotides.

Instructions for identifying DNA sequences by means of hybridization can be found by the expert, inter alia, in the handbook "The DIG System Users Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim, Germany, 1993) and in Liebl et al. (International Journal

of Systematic Bacteriology (1991) 41: 255-260). The hybridization takes place under stringent conditions, that is to say only hybrids in which the probe and target sequence, i.e. the polynucleotides treated with the probe, are at least 70% identical are formed. It is known that the stringency of the hybridization, including the washing steps, is influenced or determined by varying the buffer composition, the temperature and the salt concentration. The hybridization reaction is preferably carried out under a relatively low stringency compared with the washing steps (Hybaid Hybridisation Guide, Hybaid Limited, Teddington, UK, 1996).

A 5x SSC buffer at a temperature of approx. 50 - 68°C, for example, can be employed for the hybridization reaction. Probes can also hybridize here with polynucleotides which are less than 70% identical to the sequence of the probe. Such hybrids are less stable and are removed by washing under stringent conditions. This can be achieved, for example, by lowering the salt concentration to 2x SSC and subsequently 0.5x SSC (The DIG System User's Guide for Filter Hybridisation, Boehringer Mannheim, Mannheim, Germany, 1995) a temperature of approx. 50 - 68°C being established. It is optionally possible to lower the salt concentration to 0.1x SSC. Polynucleotide fragments which are, for example, at least 70% or at least 80% or at least 90% to 95% identical to the sequence of the probe employed can be isolated by increasing the hybridization temperature stepwise in steps of approx. 1 - 2°C. Further instructions on hybridization are obtainable on the market in the form of so-called kits (e.g. DIG Easy Hyb from Roche Diagnostics GmbH, Mannheim, Germany, Catalogue No. 1603558).

Instructions for amplification of DNA sequences with the aid of the polymerase chain reaction (PCR) can be found by the expert, inter alia, in the handbook by Gait: Oligonukleotide synthesis: A Practical Approach (IRL Press,

Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994).

It has been found that coryneform bacteria produce amino acids, in particular L-lysine, in an improved manner after
5 over-expression of the *cstA* gene.

To achieve an over-expression, the number of copies of the corresponding genes can be increased, or the promoter and regulation region or the ribosome binding site upstream of the structural gene can be mutated. Expression cassettes
10 which are incorporated upstream of the structural gene act in the same way. By inducible promoters, it is additionally possible to increase the expression in the course of fermentative lysine production. The expression is likewise improved by measures to prolong the life of the
15 m-RNA. Furthermore, the enzyme activity is also increased by preventing the degradation of the enzyme protein. The genes or gene constructs can either be present in plasmids with a varying number of copies, or can be integrated and amplified in the chromosome. Alternatively, an over-
20 expression of the genes in question can furthermore be achieved by changing the composition of the media and the culture procedure.

Instructions in this context can be found by the expert, inter alia, in Martin et al. (Bio/Technology 5, 137-146
25 (1987)), in Guerrero et al. (Gene 138, 35-41 (1994)), Tsuchiya and Morinaga (Bio/Technology 6, 428-430 (1988)), in Eikmanns et al. (Gene 102, 93-98 (1991)), in European Patent Specification 0 472 869, in US Patent 4,601,893, in Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)), in
30 Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)), in LaBarre et al. (Journal of Bacteriology 175, 1001-1007 (1993)), in Patent Application WO 96/15246, in Malumbres et al. (Gene 134, 15 - 24 (1993)), in Japanese Laid-Open Specification JP-A-10-
35 229891, in Jensen and Hammer (Biotechnology and

Bioengineering 58, 191-195 (1998)), in Makrides (Microbiological Reviews 60:512-538 (1996)) and in known textbooks of genetics and molecular biology.

By way of example, for enhancement the *cstA* gene according to the invention was over-expressed with the aid of episomal plasmids. Suitable plasmids are those which are replicated in coryneform bacteria. Numerous known plasmid vectors, such as e.g. pZ1 (Menkel et al., Applied and Environmental Microbiology (1989) 64: 549-554), pEKEx1 (Eikmanns et al., Gene 102:93-98 (1991)) or pHS2-1 (Sonnen et al., Gene 107:69-74 (1991)) are based on the cryptic plasmids pHM1519, pBL1 or pGA1. Other plasmid vectors, such as e.g. those based on pCG4 (US-A 4,489,160), or pNG2 (Serwold-Davis et al., FEMS Microbiology Letters 66, 119-124 (1990)), or pAG1 (US-A 5,158,891), can be used in the same manner.

Plasmid vectors which are furthermore suitable are also those with the aid of which the process of gene amplification by integration into the chromosome can be used, as has been described, for example, by Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)) for duplication or amplification of the *hom-thrB* operon. In this method, the complete gene is cloned in a plasmid vector which can replicate in a host (typically *E. coli*), but not in *C. glutamicum*. Possible vectors are, for example, pSUP301 (Simon et al., Bio/Technology 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-73 (1994)), pGEM-T (Promega corporation, Madison, WI, USA), pCR2.1-TOPO (Shuman (1994). Journal of Biological Chemistry 269:32678-84; US-A 5,487,993), pCR®Blunt (Invitrogen, Groningen, Holland; Bernard et al., Journal of Molecular Biology, 234: 534-541 (1993)), pEM1 (Schrumpf et al, 1991, Journal of Bacteriology 173:4510-4516) or pBGS8 (Spratt et al., 1986, Gene 41: 337-342). The plasmid vector which contains the gene to be amplified is then transferred into the desired strain of *C. glutamicum* by conjugation or

transformation. The method of conjugation is described, for example, by Schäfer et al. (Applied and Environmental Microbiology 60, 756-759 (1994)). Methods for transformation are described, for example, by Thierbach et al. (Applied Microbiology and Biotechnology 29, 356-362 (1988)), Dunican and Shivnan (Bio/Technology 7, 1067-1070 (1989)) and Tauch et al. (FEMS Microbiological Letters 123, 343-347 (1994)). After homologous recombination by means of a "cross over" event, the resulting strain contains at least two copies of the gene in question.

In addition, it may be advantageous for the production of amino acids, in particular L-lysine, to enhance one or more enzymes of the particular biosynthesis pathway, of glycolysis, of anaplerosis, of the citric acid cycle or of amino acid export and optionally regulatory proteins, in addition to the *cstA* gene.

Thus, for example, for the preparation of amino acids, in particular L-lysine, one or more genes chosen from the group consisting of

- the *dapA* gene which codes for dihydrodipicolinate synthase (EP-B 0 197 335),
- the *gap* gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the *tpi* gene which codes for triose phosphate isomerase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the *pgk* gene which codes for 3-phosphoglycerate kinase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the *pyc* gene which codes for pyruvate carboxylase (Peters-Wendisch et al. (Microbiology 144, 915 - 927 (1998)),

- the lysC gene which codes for a feed back resistant aspartate kinase (Kalinowski et al. (1990), Molecular and General Genetics 224, 317-324; Accession No.P26512),
- the lysE gene which codes for lysine export,
5 (DE-A-195 48 222),
- the zwal gene which codes for the Zwal protein (DE: 19959328.0, DSM 13115)

can be enhanced, in particular over-expressed.

10 It may furthermore be advantageous for the production of amino acids, in particular L-lysine, in addition to the enhancement of the cstA gene, for one or more genes chosen from the group consisting of

- the pck gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1; DSM 13047),
- 15 • the pgi gene which codes for glucose 6-phosphate isomerase (US 09/396,478; DSM 12969),
- the poxB gene which codes for pyruvate oxidase (DE: 1995 1975.7; DSM 13114),
- the zwa2 gene which codes for the Zwa2 protein (DE:
20 19959327.2, DSM 13113)

to be attenuated, in particular for the expression thereof to be reduced.

25 In addition to over-expression of the cstA gene it may furthermore be advantageous, for the production of amino acids, in particular L-lysine, to eliminate undesirable side reactions, (Nakayama: "Breeding of Amino Acid Producing Micro-organisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The microorganisms prepared according to the invention can be cultured continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of production of amino acids, in particular L-lysine. A summary of known culture methods are described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as e.g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as e.g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e.g. glycerol and ethanol, and organic acids, such as e.g. acetic acid, can be used as the source of carbon. These substance can be used individually or as a mixture.

Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus.

- The culture medium must furthermore comprise salts of metals, such as e. g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the abovementioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.
- Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, such as e.g. antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as e.g. air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of the desired product has formed. This target is usually reached within 10 hours to 160 hours.
- The analysis of lysine can be carried out by ion exchange chromatography with subsequent ninhydrin derivatization, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190).
- The process according to the invention is used for the fermentative preparation of amino acids, in particular L-lysine.

The present invention is explained in more detail in the following with the aid of embodiment examples.

The isolation of plasmid DNA from *Escherichia coli* and all techniques of restriction, Klenow and alkaline phosphatase treatment were carried out by the method of Sambrook et al. (Molecular Cloning. A Laboratory Manual (1989) Cold Spring Harbour Laboratory Press, Cold Spring Harbor, NY, USA). Methods for transformation of *Escherichia coli* are also described in this handbook.

The composition of the usual nutrient media, such as LB or TY medium, can also be found in the handbook by Sambrook et al.

Example 1

Preparation of a genomic cosmid gene library from *Corynebacterium glutamicum* ATCC 13032

Chromosomal DNA from *Corynebacterium glutamicum* ATCC 13032 was isolated as described by Tauch et al. (1995, Plasmid 33:168-179) and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Code no. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Code no. 1758250). The DNA of the cosmid vector SuperCos1 (Wahl et al. (1987) Proceedings of the National Academy of Sciences USA 84:2160-2164), obtained from Stratagene (La Jolla, USA, Product Description SuperCos1 Cosmid Vector Kit, Code no. 251301) was cleaved with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, Product Description XbaI, Code no. 27-0948-02) and likewise dephosphorylated with shrimp alkaline phosphatase.

The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Code no. 27-0868-04). The cosmid DNA treated in this manner was mixed with the treated ATCC13032 DNA and the batch was treated with T4 DNA ligase (Amersham

Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04). The ligation mixture was then packed in phages with the aid of Gigapack II XL Packing Extract (Stratagene, La Jolla, USA, Product Description
5 Gigapack II XL Packing Extract, Code no. 200217).

For infection of the E. coli strain NM554 (Raleigh et al. 1988, Nucleic Acid Research 16:1563-1575) the cells were taken up in 10 mM MgSO₄ and mixed with an aliquot of the phage suspension. The infection and titering of the cosmid
10 library were carried out as described by Sambrook et al. (1989, Molecular Cloning: A laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 1955, Virology, 1:190) with 100 mg/l ampicillin. After incubation overnight at 37°C, recombinant individual clones
15 were selected.

Example 2

Isolation and sequencing of the cstA gene

The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen,
20 Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Product No. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline
25 phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Product No. 1758250). After separation by gel electrophoresis, the cosmid fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen,
30 Hilden, Germany).

The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, The Netherlands, Product Description Zero Background Cloning Kit, Product No. K2500-01) was cleaved with the restriction enzyme BamHI (Amersham

Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol Letters, 123:343-7) into the E. coli strain DH5 α MCR (Grant, 1990, Proceedings of the National Academy of Sciences U.S.A., 87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50 mg/l zeocin.

The plasmid preparation of the recombinant clones was carried out with Biorobot 9600 (Product No. 900200, Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain termination method of Sanger et al. (1977, Proceedings of the National Academy of Sciences U.S.A., 74:5463-5467) with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067). The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) (Product No. A124.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).

The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 14:217-231) version 97-0. The individual sequences of the pZero1 derivatives were assembled to a continuous contig. The computer-assisted coding region analysis was prepared with the XNIP program (Staden, 1986, Nucleic Acids Research, 14:217-231).

The resulting nucleotide sequence is shown in SEQ ID No. 1. Analysis of the nucleotide sequence showed an open reading frame of 2316 base pairs, which was called the cstA gene. The cstA gene codes for a protein of 772 amino acids.

SEQUENCE PROTOCOL

<110> Degussa-Hüls AG

5 <120> New nucleotide sequences which code for the cstA gene

<130> 000173 BT

<140>

10 <141>

<160> 2

<170> PatentIn Ver. 2.1

15

<210> 1

<211> 2718

<212> DNA

<213> Corynebacterium glutamicum

20

<220>

<221> CDS

<222> (200)..(2515)

<223> cstA gene

25

<400> 1

aggatggtat aaatcatctc tcaatgttac ttttccattg ttaagaatta acaactctcg 60

gtgatttgtc gcatacccag ctgtcaaaga tccgatcatc ggcatacaga aacacccatc 120

30

tggccgaact ttcttttttc tgcattgcatt tctgcacaca gtttctgccc gctgtttctg 180

cccgctgttt ctaogcata gtg gct ttg aaa cga ccc gaa gag aaa aca gta 232

35

Val Ala Leu Lys Arg Pro Glu Glu Lys Thr Val
1 5 10

aag atc gtg acc ata aaa cag act gac aac atc aat gac gat gat ttg 280

Lys Ile Val Thr Ile Lys Gln Thr Asp Asn Ile Asn Asp Asp Asp Leu
15 20 25

40

gtg tac agc aac gct act gac ctt cca gta ggc gtg aag aag tcc cct 328

Val Tyr Ser Asn Ala Thr Asp Leu Pro Val Gly Val Lys Lys Ser Pro
30 35 40

45

aaa atg tca ccg acc gcc cgc gtt ggt ctc ctt gtc ttt ggg gtt atc 376

Lys Met Ser Pro Thr Ala Arg Val Gly Leu Leu Val Phe Gly Val Ile
45 50 55

50

gcg gcg gtg ggt tgg gga gca atc gct ttc tcc cgt ggc gaa aca atc 424

Ala Ala Val Gly Trp Gly Ala Ile Ala Phe Ser Arg Gly Glu Thr Ile
60 65 70 75

55

aac tct gtg tgg ctg gtt ttg gcg gca gtt ggt tcc tat atc att gcg 472

Asn Ser Val Trp Leu Val Leu Ala Ala Val Gly Ser Tyr Ile Ile Ala
80 85 90

	ttt tct ttc tat gcc cga ctg att gaa tac aaa gtt gtt aag ccg aaa	520
	Phe Ser Phe Tyr Ala Arg Leu Ile Glu Tyr Lys Val Val Lys Pro Lys	
	95 100 105	
5	gat cag cga gca acc ccg gcg gaa tac gtt aat gac ggc aag gac tat	568
	Asp Gln Arg Ala Thr Pro Ala Glu Tyr Val Asn Asp Gly Lys Asp Tyr	
	110 115 120	
10	gtc cca acg gat cgt cgt gtg ctt ttt ggc cac cac ttt gca gct att	616
	Val Pro Thr Asp Arg Arg Val Leu Phe Gly His His Phe Ala Ala Ile	
	125 130 135	
15	gca ggt gcc ggt cca ttg gtt gga cct gtc atg gcc gcg cag atg ggc	664
	Ala Gly Ala Gly Pro Leu Val Gly Pro Val Met Ala Ala Gln Met Gly	
	140 145 150 155	
20	tac ctg cca ggc acc ttg tgg att atc ctc ggt gtg att ttc gcc ggt	712
	Tyr Leu Pro Gly Thr Leu Trp Ile Ile Leu Gly Val Ile Phe Ala Gly	
	160 165 170	
25	gca gtg cag gac tac cta gtg ctg tgg gtg tct act cgt agg cgt gga	760
	Ala Val Gln Asp Tyr Leu Val Leu Trp Val Ser Thr Arg Arg Arg Gly	
	175 180 185	
30	cgc tca ctt ggc cag atg gtt cgt gat gaa atg ggc acg gtc ggt gga	808
	Arg Ser Leu Gly Gln Met Val Arg Asp Glu Met Gly Thr Val Gly Gly	
	190 195 200	
35	gct gcc ggt atc ttg gcg acc atc tcc atc atg atc atc att atc gcg	856
	Ala Ala Gly Ile Leu Ala Thr Ile Ser Ile Met Ile Ile Ile Ala	
	205 210 215	
40	gtg ctc gca ttg atc gtg gtt aat gca ctg gct gat tca cca tgg ggc	904
	Val Leu Ala Leu Ile Val Val Asn Ala Leu Ala Asp Ser Pro Trp Gly	
	220 225 230 235	
45	gtt ttc tcc atc acc atg acc atc cca att gca ctg ttc atg ggt gtg	952
	Val Phe Ser Ile Thr Met Thr Ile Pro Ile Ala Leu Phe Met Gly Val	
	240 245 250	
50	tac ttg cgt tac ctg cgc cca ggt cgt gtt act gaa gtg tcc atc atc	1000
	Tyr Leu Arg Tyr Leu Arg Pro Gly Arg Val Thr Glu Val Ser Ile Ile	
	255 260 265	
55	ggt gtg gca ctg ctc ctg ctg gct atc gtt gct ggt ggt tgg gtt gca	1048
	Gly Val Ala Leu Leu Leu Leu Ala Ile Val Ala Gly Gly Trp Val Ala	
	270 275 280	
60	gac acc tca tgg ggc gtg gaa tgg ttc acc tgg tct aag acc act ttg	1096
	Asp Thr Ser Trp Gly Val Glu Trp Phe Thr Trp Ser Lys Thr Thr Leu	
	285 290 295	
65	gcg ttg gcc ttg atc ggt tac gga atc atg gct gcg att ttg ccg gtg	1144
	Ala Leu Ala Leu Ile Gly Tyr Gly Ile Met Ala Ala Ile Leu Pro Val	
	300 305 310 315	
70	tgg ctg ctg ctt gca ccg cgc gat tac ctg tct acc ttt atg aag atc	1192
	Trp Leu Leu Leu Ala Pro Arg Asp Tyr Leu Ser Thr Phe Met Lys Ile	
	320 325 330	

	ggc gtc atc ggt ctg ttg gca gtg ggt att ttg ttc gca cgt cct gag	1240
	Gly Val Ile Gly Leu Leu Ala Val Gly Ile Leu Phe Ala Arg Pro Glu	
	335 340 345	
5	gtg cag atg cct tcc gtg acc tcc ttc gca ctt gag ggc aac ggt ccg	1288
	Val Gln Met Pro Ser Val Thr Ser Phe Ala Leu Glu Gly Asn Gly Pro	
	350 355 360	
10	gtg ttc tct gga agt ctg ttc cca ttc ctg ttc atc acg att gcc tgt	1336
	Val Phe Ser Gly Ser Leu Phe Pro Phe Leu Phe Ile Thr Ile Ala Cys	
	365 370 375	
15	ggt gca ctg tct ggt ttc cac gca ctg att tct tca gga acc aca cca	1384
	Gly Ala Leu Ser Gly Phe His Ala Leu Ile Ser Ser Gly Thr Thr Pro	
	380 385 390 395	
20	aag ctt gtg gag aag gaa tcc cag atg cgc atg ctc ggc tac ggc ggc	1432
	Lys Leu Val Glu Lys Glu Ser Gln Met Arg Met Leu Gly Tyr Gly Gly	
	400 405 410	
25	atg ttg atg gaa tct ttc gtg gcg atg atg gca ctg atc acc gct gtt	1480
	Met Leu Met Glu Ser Phe Val Ala Met Met Ala Leu Ile Thr Ala Val	
	415 420 425	
	att ctg gat cgt cac ctg tac ttc tcc atg aac gct ccg ctg gca ctg	1528
	Ile Leu Asp Arg His Leu Tyr Phe Ser Met Asn Ala Pro Leu Ala Leu	
	430 435 440	
30	act ggt gga gat cca gca acc gca gct gag tgg gtt aac tcc att ggg	1576
	Thr Gly Gly Asp Pro Ala Thr Ala Ala Glu Trp Val Asn Ser Ile Gly	
	445 450 455	
35	ctg aca ggt gcg gat atc acc ccg gaa cag ctg tcg gaa gct gct gaa	1624
	Leu Thr Gly Ala Asp Ile Thr Pro Glu Gln Leu Ser Glu Ala Ala Glu	
	460 465 470 475	
40	agt gtc gga gaa tcc act gtt att tcc cgt acc ggt ggc gca cca acc	1672
	Ser Val Gly Glu Ser Thr Val Ile Ser Arg Thr Gly Gly Ala Pro Thr	
	480 485 490	
45	ttg gcg ttc ggt atg tct gaa atc ctc tcc gga ttc atc ggc ggc gct	1720
	Leu Ala Phe Gly Met Ser Glu Ile Leu Ser Gly Phe Ile Gly Gly Ala	
	495 500 505	
	gga atg aag gcg ttc tgg tac cac ttc gcc atc atg ttt gag gct ctg	1768
	Gly Met Lys Ala Phe Trp Tyr His Phe Ala Ile Met Phe Glu Ala Leu	
	510 515 520	
50	ttc atc ctc act act gtg gat gca ggt act cgt gtg gct cgc ttt atg	1816
	Phe Ile Leu Thr Thr Val Asp Ala Gly Thr Arg Val Ala Arg Phe Met	
	525 530 535	
55	atg acc gat acc ttg ggc aat gtt cca ggt ctg cgc cgt ttc aag gat	1864
	Met Thr Asp Thr Leu Gly Asn Val Pro Gly Leu Arg Arg Phe Lys Asp	
	540 545 550 555	

	cct tca tgg act gtc ggt aac tgg att tct acc gtg ttt gtg tgt gct	1912
	Pro Ser Trp Thr Val Gly Asn Trp Ile Ser Thr Val Phe Val Cys Ala	
	560 565 570	
5	cta tgg ggt gct att ttg ctc atg ggt gtt acc gat cca ctg ggc ggc	1960
	Leu Trp Gly Ala Ile Leu Leu Met Gly Val Thr Asp Pro Leu Gly Gly	
	575 580 585	
10	atc aac gtg ctt ttc cca cta ttc ggt atc gct aac cag ctg ctc gcc	2008
	Ile Asn Val Leu Phe Pro Leu Phe Gly Ile Ala Asn Gln Leu Leu Ala	
	590 595 600	
15	gct att gca ctt gct ctc gtg ctg gtt gtt gtg gtg aag aag ggc ctg	2056
	Ala Ile Ala Leu Ala Leu Val Leu Val Val Val Val Lys Lys Gly Leu	
	605 610 615	
20	tac aag tgg gcg tgg att cca gct gtt cct ttg gca tgg gat ctc att	2104
	Tyr Lys Trp Ala Trp Ile Pro Ala Val Pro Leu Ala Trp Asp Leu Ile	
	620 625 630 635	
25	gtc acg atg act gcg tca tgg cag aag att ttc cac tct gat ccg gct	2152
	Val Thr Met Thr Ala Ser Trp Gln Lys Ile Phe His Ser Asp Pro Ala	
	640 645 650	
30	att ggc tac tgg gct cag aac gcg aac ttc cgc gat gca aag tct caa	2200
	Ile Gly Tyr Trp Ala Gln Asn Ala Asn Phe Arg Asp Ala Lys Ser Gln	
	655 660 665	
35	ggc ctt acc gaa ttt ggt gcc gct aaa tct cct gag gca atc gat gcg	2248
	Gly Leu Thr Glu Phe Gly Ala Ala Lys Ser Pro Glu Ala Ile Asp Ala	
	670 675 680	
40	gtt atc cga aac acc atg att cag ggc atc ttg tcc atc ctg ttc gcg	2296
	Val Ile Arg Asn Thr Met Ile Gln Gly Ile Leu Ser Ile Leu Phe Ala	
	685 690 695	
45	gtg ctc gtc ctc gtt gtt gtc ggc gca gcc att gcg gtg tgc atc aag	2344
	Val Leu Val Leu Val Val Val Gly Ala Ala Ile Ala Val Cys Ile Lys	
	700 705 710 715	
50	tcc atc agg gct cgt gca gcc gga aca cct ttg gag acc act gaa gag	2392
	Ser Ile Arg Ala Arg Ala Ala Gly Thr Pro Leu Glu Thr Thr Glu Glu	
	720 725 730	
55	cct gat act gaa tct gag ttc ttc gcc cca act gga ttc ctt gca tct	2440
	Pro Asp Thr Glu Ser Glu Phe Phe Ala Pro Thr Gly Phe Leu Ala Ser	
	735 740 745	
60	tcc agg gat aag gaa gtc cag gcc atg tgg gac gag cgc tac cca ggc	2488
	Ser Arg Asp Lys Glu Val Gln Ala Met Trp Asp Glu Arg Tyr Pro Gly	
	750 755 760	
65	ggg ggc ccc gtg tct tct gga ggg cac taaaacatga tggctcttac	2535
	Gly Ala Pro Val Ser Ser Gly Gly His	
	765 770	
	tcatgcactg tggaaaatcc cgcgggcggt gtggtggtat ctactgagc tcatggggga	2595
	cacggcggtat tccaagtatg tggtgcactt aaagcaccac catccggatg ctccgattcc	2655

tactgagcgg gagtattggc gggcaaagta tgcagatcag gacgctaatac ctggtgccccg 2715
 ctg 2718

5

<210> 2
 <211> 772
 <212> PRT
 10 <213> Corynebacterium glutamicum

<400> 2
 Val Ala Leu Lys Arg Pro Glu Glu Lys Thr Val Lys Ile Val Thr Ile
 1 5 10 15
 15 Lys Gln Thr Asp Asn Ile Asn Asp Asp Asp Leu Val Tyr Ser Asn Ala
 20 25 30
 20 Thr Asp Leu Pro Val Gly Val Lys Lys Ser Pro Lys Met Ser Pro Thr
 35 40 45
 Ala Arg Val Gly Leu Leu Val Phe Gly Val Ile Ala Ala Val Gly Trp
 50 55 60
 25 Gly Ala Ile Ala Phe Ser Arg Gly Glu Thr Ile Asn Ser Val Trp Leu
 65 70 75 80
 Val Leu Ala Ala Val Gly Ser Tyr Ile Ile Ala Phe Ser Phe Tyr Ala
 85 90 95
 30 Arg Leu Ile Glu Tyr Lys Val Val Lys Pro Lys Asp Gln Arg Ala Thr
 100 105 110
 Pro Ala Glu Tyr Val Asn Asp Gly Lys Asp Tyr Val Pro Thr Asp Arg
 115 120 125
 Arg Val Leu Phe Gly His His Phe Ala Ala Ile Ala Gly Ala Gly Pro
 130 135 140
 40 Leu Val Gly Pro Val Met Ala Ala Gln Met Gly Tyr Leu Pro Gly Thr
 145 150 155 160
 Leu Trp Ile Ile Leu Gly Val Ile Phe Ala Gly Ala Val Gln Asp Tyr
 165 170 175
 45 Leu Val Leu Trp Val Ser Thr Arg Arg Arg Gly Arg Ser Leu Gly Gln
 180 185 190
 Met Val Arg Asp Glu Met Gly Thr Val Gly Gly Ala Ala Gly Ile Leu
 195 200 205
 Ala Thr Ile Ser Ile Met Ile Ile Ile Ile Ala Val Leu Ala Leu Ile
 210 215 220
 55 Val Val Asn Ala Leu Ala Asp Ser Pro Trp Gly Val Phe Ser Ile Thr
 225 230 235 240
 Met Thr Ile Pro Ile Ala Leu Phe Met Gly Val Tyr Leu Arg Tyr Leu
 245 250 255

	Arg	Pro	Gly	Arg	Val	Thr	Glu	Val	Ser	Ile	Ile	Gly	Val	Ala	Leu	Leu	
				260					265					270			
5	Leu	Leu	Ala	Ile	Val	Ala	Gly	Gly	Trp	Val	Ala	Asp	Thr	Ser	Trp	Gly	
			275					280					285				
	Val	Glu	Trp	Phe	Thr	Trp	Ser	Lys	Thr	Thr	Leu	Ala	Leu	Ala	Leu	Ile	
		290					295					300					
10	Gly	Tyr	Gly	Ile	Met	Ala	Ala	Ile	Leu	Pro	Val	Trp	Leu	Leu	Leu	Ala	
	305					310					315					320	
	Pro	Arg	Asp	Tyr	Leu	Ser	Thr	Phe	Met	Lys	Ile	Gly	Val	Ile	Gly	Leu	
15					325					330					335		
	Leu	Ala	Val	Gly	Ile	Leu	Phe	Ala	Arg	Pro	Glu	Val	Gln	Met	Pro	Ser	
				340					345					350			
20	Val	Thr	Ser	Phe	Ala	Leu	Glu	Gly	Asn	Gly	Pro	Val	Phe	Ser	Gly	Ser	
			355					360					365				
	Leu	Phe	Pro	Phe	Leu	Phe	Ile	Thr	Ile	Ala	Cys	Gly	Ala	Leu	Ser	Gly	
		370					375					380					
25	Phe	His	Ala	Leu	Ile	Ser	Ser	Gly	Thr	Thr	Pro	Lys	Leu	Val	Glu	Lys	
	385					390					395					400	
	Glu	Ser	Gln	Met	Arg	Met	Leu	Gly	Tyr	Gly	Gly	Met	Leu	Met	Glu	Ser	
30					405					410					415		
	Phe	Val	Ala	Met	Met	Ala	Leu	Ile	Thr	Ala	Val	Ile	Leu	Asp	Arg	His	
				420					425					430			
35	Leu	Tyr	Phe	Ser	Met	Asn	Ala	Pro	Leu	Ala	Leu	Thr	Gly	Gly	Asp	Pro	
			435					440					445				
	Ala	Thr	Ala	Ala	Glu	Trp	Val	Asn	Ser	Ile	Gly	Leu	Thr	Gly	Ala	Asp	
		450					455					460					
40	Ile	Thr	Pro	Glu	Gln	Leu	Ser	Glu	Ala	Ala	Glu	Ser	Val	Gly	Glu	Ser	
	465					470					475					480	
	Thr	Val	Ile	Ser	Arg	Thr	Gly	Gly	Ala	Pro	Thr	Leu	Ala	Phe	Gly	Met	
45					485					490					495		
	Ser	Glu	Ile	Leu	Ser	Gly	Phe	Ile	Gly	Gly	Ala	Gly	Met	Lys	Ala	Phe	
				500					505					510			
50	Trp	Tyr	His	Phe	Ala	Ile	Met	Phe	Glu	Ala	Leu	Phe	Ile	Leu	Thr	Thr	
			515					520					525				
	Val	Asp	Ala	Gly	Thr	Arg	Val	Ala	Arg	Phe	Met	Met	Thr	Asp	Thr	Leu	
		530					535					540					
55	Gly	Asn	Val	Pro	Gly	Leu	Arg	Arg	Phe	Lys	Asp	Pro	Ser	Trp	Thr	Val	
	545					550					555					560	

[illegible]

Patent claims

1. An isolated polynucleotide from coryneform bacteria,
comprising a polynucleotide sequence which codes for
5 the *cstA* gene, chosen from the group consisting of
 - a) polynucleotide which is identical to the extent of at
least 70% to a polynucleotide which codes for a
polypeptide which comprises the amino acid sequence
of SEQ ID No. 2,
 - 10 b) polynucleotide which codes for a polypeptide which
comprises an amino acid sequence which is identical
to the extent of at least 70% to the amino acid
sequence of SEQ ID No. 2,
 - c) polynucleotide which is complementary to the
15 polynucleotides of a) or b), and
 - d) polynucleotide comprising at least 15 successive
nucleotides of the polynucleotide sequence of a), b)
or c),

the polypeptide preferably having the activity of carbon
20 starvation protein A.
2. The polynucleotide as claimed in claim 1, wherein the
polynucleotide is a preferably recombinant DNA which is
capable of replication in coryneform bacteria.
3. The polynucleotide as claimed in claim 1, wherein the
25 polynucleotide is an RNA.
4. The polynucleotide as claimed in claim 2, comprising
the nucleic acid sequence as shown in SEQ ID No. 1.
5. The DNA as claimed in claim 2 which is capable of
replication, comprising
30 (i) the nucleotide sequence shown in SEQ ID No. 1, or

- (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
 - (iii) at least one sequence which hybridizes with the
5 sequence complementary to sequence (i) or (ii), and optionally
 - (iv) sense mutations of neutral function in (i).
6. The polynucleotide sequence as claimed in claim 2,
which codes for a polypeptide which comprises the amino
10 acid sequences shown in SEQ ID No. 2.
 7. Coryneform bacteria in which the cstA gene is enhanced,
in particular over-expressed.
 8. A process for the fermentative preparation of L-amino
acids, in particular L-lysine, wherein the following
15 steps are carried out:
 - a) fermentation of the coryneform bacteria which produce
the desired L-amino acid and in which at least the
cstA gene or nucleotide sequences which code for it
are enhanced, in particular over-expressed;
 - 20 b) concentration of the L-amino acid in the medium or in
the cells of the bacteria, and
 - c) isolation of the L-amino acid.
 9. The process as claimed in claim 8, wherein bacteria in
which further genes of the biosynthesis pathway of the
25 desired L-amino acid are additionally enhanced are
employed.
 10. The process as claimed in claim 8, wherein bacteria in
which the metabolic pathways which reduce the formation
of the desired L-amino acid are at least partly
30 eliminated are employed.

11. The process as claimed in claim 8, wherein a strain transformed with a plasmid vector is employed, and the plasmid vector carries the nucleotide sequence which codes for the *cstA* gene.
- 5 12. The process as claimed in claim 8, wherein the expression of the polynucleotide which codes for the *cstA* gene is enhanced, in particular over-expressed.
13. The process as claimed in claim 8, wherein the regulatory properties of the polypeptide for which the
10 polynucleotide *cstA* codes are increased.
14. The process as claimed in claim 8, wherein for the preparation of L-amino acids, in particular L-lysine, coryneform microorganisms in which at the same time one or more of the genes chosen from the group consisting
15 of
- 14.1 the *lysC* gene which codes for a feed back resistant aspartate kinase,
- 14.2 the *dapA* gene which codes for dihydrodipicolinate synthase,
- 20 14.3 the *gap* gene which codes for glycerolaldehyde 3-phosphate dehydrogenase,
- 14.4 the *pgk* gene which codes for 3-phosphoglycerate kinase,
- 14.5 the *pyc* gene which codes for pyruvate
25 carboxylase,
- 14.6 the *tpi* gene which codes for triose phosphate isomerase,
- 14.7 the *lysE* gene which codes for lysine export,
- 14.8 the *zwf* gene which codes for the *Zwf* protein,

is or are enhanced or over-expressed are fermented.

15. The process as claimed in claim 8, wherein for the preparation of L-amino acids, in particular L-lysine, coryneform microorganisms in which at the same time one or more of the genes chosen from the group consisting of

15.1 the pck gene which codes for phosphoenolpyruvate carboxykinase,

15.2 the pgi gene which codes for glucose 6-phosphate6 isomerase,

15.3 the poxB gene which codes for pyruvate oxidase,

15.4 the zwa2 gene which codes for the Zwa2 protein

is or are attenuated are fermented.

16. Coryneform bacteria which contain a vector which carries a polynucleotide as claimed in claim 1.

17. The process as claimed in one or more of the preceding claims, wherein microorganisms of the genus *Corynebacterium glutamicum* are employed.

18. A process for discovering RNA, cDNA and DNA in order to isolate nucleic acids, or polynucleotides or genes which code for carbon starvation protein A or have a high similarity with the sequence of the cstA gene, wherein the polynucleotide sequences as claimed in claim 1, 2, 3 or 4 are employed as hybridization probes.

19. The process as claimed in claim 18, wherein the hybridization is carried out under a stringency corresponding to at most 2x SSC.

New nucleotide sequences which code for the cstA gene

Abstract

The invention relates to an isolated polynucleotide comprising a polynucleotide sequence chosen from the group
5 consisting of

- a) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2,
- 10 b) polynucleotide which codes for a polypeptide which comprises an amino acid sequence which is identical to the extent of at least 70% to the amino acid sequence of SEQ ID No. 2,
- c) polynucleotide which is complementary to the
15 polynucleotides of a) or b), and
- d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequence of a), b) or c),

and a process for the fermentative preparation of L-amino
20 acids using coryneform bacteria in which at least the cstA gene is present in enhanced form, and the use of the polynucleotide sequences as hybridization probes.